

Rotating pasture for pregnant sows

Problem

Feed is the biggest cost in pig farming, especially in free-range systems. In these systems, feed consumption is higher due to higher activity and higher thermal regulation needs. Outdoor sows are usually housed in huts on grass leys.

Solution

During periods of high grass growth, it is possible to reduce the amount of feed and protein content given to sows in order to get the most out of grazing. The concentrated feed can be limited to 80 % of the recommended amount for at least 3 months. At the same time, this "pasture feed" can be lower in protein. Note: from 3 weeks before farrowing, sows need a complete diet.

Benefits

With this practice, feed and production costs can be reduced by 16% due to lower quantities of concentrate feed and lower costs of "pasture feed" per ton.

Applicability box

Theme

Pigs - Animal husbandry - Feed and nutrition - Production systems - Ration planning

Geographical coverage

More efficient in areas of high grassland productivity

Application time

Pasturing season

Period of impact

Pasturing season

Equipment

Movable fences to contain/move sows to desired areas; pig feeders

Best in


On-farm feed processing (possibility to process low-protein feed)

Practical Recommendations

- Allow time for the grass ley to establish before sows start grazing.
- Enrich the ley with legumes to provide good nutritional value and palatability for sows.
- Harvest/remove less palatable plants that are not eaten by sows.



Sows before accessing a new paddock. Photo: ITAB



Ration planning (concentrate feed and pasturing) for pregnant sows. Graph: ITAB

Further information

Video

- Have a look at the following video for further instructions (French): [Paturage des truies aux trinottières.](#)

Weblinks

- Check the Organic Farm Knowledge platform for more [practical recommendations on animal husbandry.](#)
- ITAB (2019, online): [Alimentation des monogastriques en agriculture biologique.](#)

About this practice abstract and OK-Net EcoFeed**Publishers:**

Research Institute of Organic Agriculture (FiBL), CH 5070 Frick,
Phone +41 62 865 72 72, info.suisse@fibl.org, www.fibl.org

IFOAM Organics Europe, BE 1000 Brussels,
Phone +32 2 280 12 23, www.organicseurope.bio,
www.organicseurope.bio

Institut de l'agriculture et de l'alimentation biologiques (ITAB), FR
75595 Paris,
Phone +33 01 40 04 50 64, , www.itab.asso.fr

Authors: Antoine Roinsard

Contact: antoine.roinsard@itab.asso.fr

Permalink:

<https://organic-farmknowledge.org/tool/35449>

**OK-Net EcoFeed:**

<https://orgprints.org/view/projects/OKNetEcoFeed.html>

This practice abstract was elaborated in the Organic Knowledge Network on Monogastric Animal Feed project. The project is running from January 2018 to December 2020. The overall aim of OKNet EcoFeed is to help farmers, breeders and the organic feed processing industry in achieving the goal of 100% use of organic and regional feed for monogastrics.

Project website: <https://ok-net-ecofeed.eu/>

Project partners:

IFOAM Organics Europe (project coordinator), BE; Aarhus University (ICROFS), DK; Organic Research Centre (ORC), UK; Institut Technique de l'Agriculture Biologique (ITAB), FR; Research Institute of Organic Agriculture (FiBL), CH; Bioland, DE; Associazione Italiana per l'Agricoltura Biologica (AIAB), IT; Donau Soja DS, AT; Swedish University of Agricultural Sciences, SE; ECOVALIA, ES; Soil Association, UK.

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 773911. This communication only reflects the author's view. The Research Executive Agency is not responsible for any use that may be made of the information provided. The authors and editors do not assume responsibility or liability for any possible factual inaccuracies or damage resulting from the application of the recommendations in this practice abstract

