

Silage feeding for laying hens

Problem

Feeding silage to organic laying hens is practised on a daily basis by many egg producers. Since hens can eat large amounts of silage, it is important to use high-quality silage.

Solution

Different types of silage can be used. However, e.g. grass, clover grass, alfalfa, hemp, horse beans, lupines and sunflower silage often have a higher protein content than e.g. barley/pea or maize silage (whole plant or the cobs only). A chemical analysis of the harvested silage is important in order to estimate the quality.

Benefits

High-quality silage can provide nutrients for the hen and encourage natural foraging behaviour, thus reducing the risk for feather pecking. Growing protein-rich silage helps increase self-sufficiency and improve the crop rotation of the organic poultry farms.

Applicability box

Theme

Animal husbandry - Feed and nutrition - Production systems - Poultry - Ration planning

Geographical coverage

Global

Application time

Feeding silage on a daily basis all year. The amount given depends on hen age and silage type.

Period of impact

During the entire laying period.

Equipment

Silage chopper, automatic system (robot) to feed the silage 1-3 times per day in the barn.

Best in

Choice of silage will depend on the soil type and if the land is often dry or waterlogged. The silages should be harvested as whole crops.

Practical Recommendations

- Choose the silage type(s) that provides the best yield and quality, depending on soil type and weather conditions.
- The silage has to be finely chopped.
- Silage generally has to be preserved under proper conditions to optimize the fermentation process.
- Analyse the protein content of the silage and other nutrients if possible.
- Avoid silage with anti-nutritional factors.
- Investing in an automatic feeding system (robot) is recommended for larger flocks to distribute the silage evenly and encourage the hens to eat it.
- Including the chemical content of the silage in the feed formulation could be an advantage, when feeding more than 20 g silage (wet weight) per hen per day.



Figure 1: Laying hens on a veranda feeding silage distributed by an automatic system (robot). Photo: Sanna Steinfeldt, AU



Figure 2: Barley-pea silage for laying hens. Photo: Niels Finn Johansen, SEGES

Further information

Weblinks

- [This video](#) provides further instructions on feeding silage to layers (in Danish).
- On the website of [Økologisk Landsforening 2019](#), there is an overview of the most important feed materials for self-supply of organic poultry with dry feed and silage. You can download and print listings of data on each crop and information on cultivation, crop rotation, harvesting, storage, nutrient content and nutritional considerations (in Danish).
- Check the Organic Farm Knowledge platform for more [practical recommendation on animal husbandry](#).

About this practice abstract and OK-Net EcoFeed**Publishers:**

Aarhus University, Department of Agroecology - Agricultural Systems and Sustainability, DK 8830 Tjele,
Phone , , <http://agro.au.dk/en>

Research Institute of Organic Agriculture (FiBL), CH 5070 Frick,
Phone +41 62 865 72 72, info.suisse@fibl.org, www.fibl.org

IFOAM Organics Europe, BE 1000 Brussels,
Phone +32 2 280 12 23, www.organicseurope.bio,
www.organicseurope.bio

Authors: Sanna Steinfeldt (ANIS-AU)

Contact: sanna.steenfeldt@anis.au.dk

Permalink:

<https://organic-farmknowledge.org/tool/35470>

**OK-Net EcoFeed:**

<https://orgprints.org/view/projects/OKNetEcoFeed.html>

This practice abstract was elaborated in the Organic Knowledge Network on Monogastric Animal Feed project. The project is running from January 2018 to December 2020. The overall aim of OKNet EcoFeed is to help farmers, breeders and the organic feed processing industry in achieving the goal of 100% use of organic and regional feed for monogastrics.

Project website: <https://ok-net-ecofeed.eu/>

Project partners:

IFOAM Organics Europe (project coordinator), BE; Aarhus University (ICROFS), DK; Organic Research Centre (ORC), UK; Institut Technique de l'Agriculture Biologique (ITAB), FR; Research Institute of Organic Agriculture (FiBL), CH; Bioland, DE; Associazione Italiana per l'Agricoltura Biologica (AIAB), IT; Donau Soja DS, AT; Swedish University of Agricultural Sciences, SE; ECOVALIA, ES; Soil Association, UK.

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 773911. This communication only reflects the author's view. The Research Executive Agency is not responsible for any use that may be made of the information provided. The authors and editors do not assume responsibility or liability for any possible factual inaccuracies or damage resulting from the application of the recommendations in this practice abstract

